

Classification of the Family Sepsidae from Korea (Diptera: Acalypttratae)

Sang Jae SUH¹ and Yong Jung KWON²

¹Department of Herb & Food Science, Tongkuk Junior College, Chilgok 718-850, Korea

²Department of Agricultural Biology, Kyungpook National University, Taegu 702-701, Korea

Abstract A total of 12 species belonging to 8 genera of Korean sepsid flies are treated herein. Among them, 2 species with 2 genera are newly added to Korean fauna. Keys and distributional data are given for each species.

Key words Classification, Diptera, Cyclorrapha, Acalypttratae, Sepsidae, Korea.

INTRODUCTION

The sepsids are small and shining blackish flies which characterized generally by having the fore femora with stout bristles or spines on the ventral surface in the males. The larvae are saprophagous and have been found in excrement, dung, manure heap, and other various decaying materials. The adults are common flies, and are often sought in considerable numbers near materials in which the larvae breed.

The first record of this family from Korea was given by Lee and Kwon (1981) who reported *Sepsis monostigma* during their faunistic investigation to Ullungdo. Then, 10 species belonging to 6 genera have been known in Korea prior to the present survey. This time, we intend to add 2 previously unrecorded species to Korean fauna. As the result, a total of 12 species belonging to 8 genera are listed here.

SYSTEMATICS

Key to the Genera of Korean Sepsidae

1. First and second basal cells united *Saltella*
– First and second basal cells separated 2
2. Mesopleural and outer vertical setae absent *Themira*
– Mesopleural and outer vertical setae present 3
3. Humeral seta absent *Decachaetophora*
– Humeral seta present 4
4. Fronto-orbital seta strong 5




Fig. 1. Fore femur of male, anterior view.

1: *Saltella orientalis*, 2: *Decachaetophora aeneipes*, 3: *Dicranosepsis bicolor*, 4: *Nemopoda pectinulata*, 5: *Sepsis bicornuta*, 6: *S. indica*, 7: *S. monostigma*, 8: *S. punctum*, 9: *S. thoracica*, 10: *Themira nigricornis*, 11: *Xenosepsis fukuharai* (scale: 0.25 mm).

7. Hypopygial process bifurcate at apex *Dicranosepsis*
 – Hypopygial process not bifurcate at apex *Sepsis*

Genus 1. *Saltella* Robineau-Desvoidy, 1830 큰꼭지파리속(신칭)

1. *Saltella orientalis* (Hendel, 1934) 큰꼭지파리(신칭) (Figs. 1-2: 1)

Pandora orientalis Hendel, 1934, Ark. Zool. 25 A 21: 4.

Saltella orientalis: Iwasa et al., 1994, Esakia 24: 204¹.

Locality. GB: Taegu, GG: Kwangŭng¹.

Distribution. Korea (Central, South), China (Gansu, Manchuria), Russia (Maritime Territory).




Fig. 2. Fore tibia of male, anterior view.

1: *Saltella orientalis*, 2: *Decachaetophora aeneipes*, 3: *Dicranosepsis bicolor*, 4: *Nemopoda pectinulata*, 5: *Sepsis bicornuta*, 6: *S. indica*, 7: *S. monostigma*, 8: *S. punctum*, 9: *S. thoracica*, 10: *Themira nigricornis*, 11: *Xenosepsis fukuharai* (scale: 0.25 mm).

Genus 2. *Decachaetophora* Duda, 1926 갈고리꼭지파리속(신칭)

2. *Decachaetophora aeneipes* (De Meijere, 1911) 갈고리꼭지파리(신칭) (Figs. 1-2: 2)

Sepsis aeneipes De Meijere, 1911, Ann. Mus. Nat. Hung. 11: 119.

Decachaetophora aeneipes: Zuska & Pont, 1984, Cat. Pal. Dipt. 9: 155 (Korea).

Locality. GN: Chirisan.

Distribution. Korea (South), Japan (Hokkaido, Honshu, Kyushu, Shikoku), China (Gansu, Sichuan, Taiwan), Russia (Maritime Territory, Kurils), Mongolia, India, Nepal, Sri Lanka, N. America.

Genus 3. *Dicranosepsis* Duda, 1926 아기꼭지파리속(신칭)

3. *Dicranosepsis bicolor* (Wiedemann, 1830) 아기꼭지파리(신칭) (Figs. 1-2: 3)

Sepsis bicolor Wiedemann, 1830, Auss. Zweifl. Ins. 2: 468.

Dicranosepsis bicolor: Iwasa et al., 1994, Esakia 24: 207-208¹⁻².

Locality. GB: Hwanghaksan, GG: Kwangnŭng¹, JB: Chŏngnŭngch'i (=Jeonglyeongchi²).

Distribution. Korea (Central, South), Japan (whole), China (Jiangsu, Taiwan), India, Indonesia, Malaysia, Nepal, Philippines, Singapore, Thailand, Vietnam.

Genus 4. *Nemopoda* Robineau-Desvoidy, 1830 산꼭지파리속(신칭)

4. *Nemopoda pectinulata* Loew, 1873 산꼭지파리(신칭) (Figs. 1-2: 4)

Nemopoda pectinulata Loew, 1873, Besch. Eur. Dipt. 3: 305.

Locality. GW: Odaesan (3ex, 19, VIII, 1986, Y. Kwon).

Distribution. Korea (new record: Central), Japan (Hokkaido, Honshu, Shikoku, Kyushu), China (Jiangsu), Russia (European part, Maritime Territory), Finland, Germany, Nepal, Norway, Sweden.

Genus 5. *Sepsis* Fallén, 1810 꼭지파리속(신칭)

Key to the species of Korean *Sepsis*

1. Pteropleuron pruinose *indica*
 – Pteropleuron shining 2
2. Stenopleuron anteroventrally shining *thoracica*
 – Stenopleuron wholly pruinose 3
3. Propleuron brown *bicornuta*
 – Propleuron black 4
4. Fore femur with a strong anterobasal seta being much longer than width of the femur
 *monostigma*
 – Fore femur with a anterobasal seta being as long as width of the femur *monostigma*

5. *Sepsis bicornuta* Ozerov, 1985 극동꼭지파리(신칭) (Figs. 1-2: 5)

Sepsis bicornuta Ozerov, 1985, Ent. Ob. 64(4): 841.

Sepsis bicornuta: Iwasa et al., 1994, Esakia 24: 205-207¹⁻³.

Locality. GB: Hwanghaksan, Sobaeksan, GG: Kwangnŭng¹, GN: Chirisan (=Tam Ri²), GW: Tong-myŏn (=Dong-myŏn³).

Distribution. Korea (Central, South), Russia (Maritime Territory).

6. *Sepsis indica* Wiedemann, 1824 노랑꼭지파리(신칭) (Figs. 1-2: 6)

Sepsis indica Wiedemann, 1824, An. Ent.: 57.

Sepsis indica: Iwasa et al., 1994, Esakia 24: 207¹.

Locality. GB: Chuwangsan, GN: Chirisan, GW: Pukpang-myŏn (=Pupan-myŏn¹).

Distribution. Korea (Central, South), Japan (whole), China (Hubei, Sichuan, Taiwan), Russia (Maritime Territory), Myanmar, India, Indonesia, Malaysia, Nepal, Philippines, Sri Lanka, Vietnam.

7. *Sepsis monostigma* Thomson, 1870 꼭지파리 (Figs. 1-2: 7)

Sepsis monostigma Thomson, 1870, Eug. Resa: 586.

Sepsis monostigma: Lee et Kwon, 1981, Rep. Kor. Ass. Cons. Nat. 19: 163¹.

Sepsis monostigma: Kim, 1984, Cheju Nat. Univ. Journ. 18: 210².

Sepsis monostigma: Kim et Chang, 1987, Rep. Kor. Ass. Cons. Nat. 25: 118³.

Sepsis monostigma: Iwasa *et al.*, 1994, Esakia 24: 205⁴⁻⁹.

Locality. GB: Chuwangsan, Hayang, Hwanghaksan, Kach'ang, Koryŏng, Kyŏngju, P'algongsan, Taegu, Ullŭngdo (=Is. Ulreung¹), GG: Kanghwado, Kwangnŭng⁴, GN: Chirisan (=Samjeong Ri⁵, Tam Ri⁶), Kajisan, Kŭmjŏngsan, GW: Obongsan, Odaesan, Sŏraksan, Taeamsan, T'aebaeksan (=Mt. T'aebaek³), Tong-myŏn (=Dong-myon⁷), Tongsan-myŏn (=Dongsan-myon⁸), JB: Maisan, Chyŏngnyŏngch'i (=Jeonglyeongchi⁹), JJ: Ilch'ulbong², JN: Tolsando, Wando.

Distribution. Korea (Central, South, Chejudo, Ullŭngdo), Japan (whole), China (Guangdong, Taiwan), Russia (Maritime Territory, Kurils), India, Philippines, Sri Lanka.

8. *Sepsis punctum* (Fabricius, 1794) 어라꼭지파리(신칭) (Figs. 1-2: 8)

Musca punctum Fabricius, 1794, Ent. Syst.: 351.

Sepsis punctum: Iwasa *et al.*, 1994, Esakia 24: 205¹⁻².

Locality. GB: P'algongsan, Sobaeksan, GN: Chirisan (=Samjeong Ri¹), GW: Obongsan, Odaesan, JB: Chŏngnyŏngch'i (=Jeonglyeongchi²), JJ: Hallasan.

Distribution. Korea (Central, South, Chejudo), Japan (Hokkaido, Honshu, Kyushu), China (Manchuria, Taiwan), Russia (European part, Kurils), Europe, N. America.

9. *Sepsis thoracica* (Robineau-Desvoidy, 1830) 등무늬꼭지파리(신칭) (Figs. 1-2: 9)

Micropeza thoracica Robineau-Desvoidy, 1830, Ess. Myod.: 742.

Sepsis thoracica: Iwasa *et al.*, 1994, Esakia 24: 207¹⁻³.

Locality. GB: Kach'ang, JN: Chindo, Chŏpto, JB: Chyŏngnyŏngch'i (=Jeonglyeongchi²), Sannae-myŏn (=Nae-lyeong Ri³).

Distribution. Korea (South), Japan (whole), China (Guangdong, Manchuria, Taiwan), India, Madagascar, Sri Lanka, Syria, Turkey, Europe, N. Africa.

Genus 6. *Themira* Robineau-Desvoidy, 1830 가시꼭지파리속(신칭)

10. *Themira nigricornis* Meigen, 1826 가시꼭지파리(신칭) (Figs. 1-2: 10)

Themira nigricornis Meigen, 1826, Syst. Bensch. 5: 291.

Locality. GB: P'algongsan (22ex, 14, VII, 1984, Y. Kwon), GN: Masan (3ex, 23, III, 1986, Y. Kwon).

Distribution. Korea (new record: South), Japan (Honshu), Russia (E. Siberia, European part), Finland, France, Germany, Kirghiz, Latvia, Norway, Sweden, UK, Ukraine.

Genus 7. *Meroplus* Rondani, 1874 애꼭지파리속(신칭)

11. *Meroplus minutus* (Wiedemann, 1830) 애꼭지파리(신칭)

Sepsis minutus Wiedemann, 1810, Auss. Zweifl. Ins. 2: 468.

Meroplus minutus: Iwasa *et al.*, 1994, Esakia 24: 207¹.

Locality. JB: Chŏngnyŏngch'i (=Jeonglyeongchi¹).

Distribution. Korea (South), Japan (Hokkaido, Honshu, Shikoku, Kyushu), China, Russia (European part, Maritime Territory), Europe, North Africa, Holarctic.

Genus 8. *Xenosepsis* Malloch, 1925 뿔꼭지파리속(신칭)

12. *Xenosepsis fukuharai* Iwasa, 1984 뿔꼭지파리(신칭) (Figs. 1-2: 11)

Xenosepsis fukuharai Iwasa, 1984, Kontyû 52: 300-302.

Xenosepsis fukuharai: Iwasa *et al.*, 1994, Esakia 24: 207¹⁻².

Locality. GB: Taegu, GW: Tongsan-myŏn (=Dongsan-myon¹), JB: Chŏngnyŏngch'i (=Jeonglyeongchi²), JN: Chindo.

Distribution. Korea (Central, South), Japan (Hokkaido, Honshu, Kyushu), China (Manchuria), Russia (Amur).

REFERENCES

- Hennig, W., 1949. Sepsidae. *Flieg. Pal. Reg.* 39a: 1-91.
- Iwasa, M., 1984. Studies on the Sepsidae from Japan (Diptera), III. On the eleven species of eight genera excluding the genera *Sepsis* Fallen and *Themira* R.-D., with description of a new species. *Kontyû* 52(2): 296-308.
- Iwasa, M., K. Kanmiya & C. E. Lee, 1994. Notes on the Sepsidae (Diptera) from South Korea. *Esakia* 34: 203-208.
- Kim, J. I. & K. S. Chang, 1987. Insect fauna of the Mt. T'aebaek in Korea. *Rep. Kor. Ass. Cons. Nat.* 25: 91-120.
- Kim, W. T., 1984. Insect fauna of four craters in Cheju Island. *Cheju Nat. Univ. Journ.* 18: 197-211.
- Lee, C. E. & Y. J. Kwon, 1981. On the insect fauna of Ulreung Is. and Dogdo Is. in Korea. *Rep. Kor. Ass. Cons. Nat.* 19: 139-178.
- Ozerov, A. L., 1985. New and little known species of the Sepsidae (Diptera) from the Far East. *Ent. Ob.* 64(4): 839-844.
- Zuska, J. & A. C. Pont, 1984. Family Sepsidae. *Cat. Pal. Dip.* 9: 154-167.

韓國産 꼭지파리科의 分類(파리목: 무판류)

서상재¹ · 권용정²

¹동국전문대학 약용식품과

²경북대학교 농과대학 농생물학과

한국산 꼭지파리류를 분류 동정한 결과, *Nemopoda pectinulata* Loew, 1873(산꼭지파리)와 *Themira nigricornis* Meigen, 1826(가시꼭지파리) 등 한국미기록 2속 2종을 새로 발견하였으며, 모두 8속 12종이 분포하고 있는 것으로 조사되었다. 또한 각 분류군의 검색표, 채집지 및 분포 등을 정리하였다.

검색어: 분류, 파리목, 무판류, 꼭지파리과

(Received: Dec. 30, 1993)

(Accepted: May 20, 1994)